

# Welcome to the **HelloHYQVIA** Free Trial Program

**HyQvia**

[Immune Globulin Infusion 10% (Human)  
with Recombinant Human Hyaluronidase]



## Infuse monthly.\* Live daily.

\*Every 3 or 4 weeks.

### What is HyQvia® [Immune Globulin Infusion 10% (Human) with Recombinant Human Hyaluronidase] Solution?

HyQvia is a liquid medicine that is given under the skin (subcutaneously) to treat primary immunodeficiency (PI) in people 2 years and older.

#### IMPORTANT SAFETY INFORMATION

##### **What is the most important information that I should know about HyQvia?**

- HyQvia can cause blood clots.
- Call your healthcare professional (HCP) if you have pain, swelling, warmth, redness, or a lump in your legs or arms, other than at the infusion site(s), unexplained shortness of breath, chest pain or discomfort that worsens on deep breathing, unexplained rapid pulse, numbness or weakness on one side of the body.
- Your HCP may perform blood tests regularly to check your IgG level.
- Do not infuse HyQvia into or around an infected or red swollen area because it can cause infection to spread.

***Please see additional safety information throughout, [click here](#) for Information for Patients, and discuss with your HCP.***

# HelloHYQVIA Free Trial Program

**HyQvia**  
[Immune Globulin Infusion 10% (Human)  
with Recombinant Human Hyaluronidase]

Here's what to expect from the Free Trial Program:



## Enroll

Once you and your healthcare provider (HCP) decide to try HyQvia and enroll in the HelloHYQVIA Free Trial Program, your HCP will start the enrollment process.

To be eligible for the HelloHYQVIA program, you must:

1. Be a patient with an ICD-10-verified diagnosis of primary immunodeficiency (PI).
2. Be a new patient not currently using HyQvia and who has not previously enrolled in the HelloHYQVIA program. Additional terms apply.

### What you need to do:

*If you do not have epinephrine at home already, you will need to get a prescription from your HCP. As a safety measure, epinephrine needs to be on hand for your first free trial HyQvia infusion.*



## Welcome call

**You will receive a welcome call from a HelloHYQVIA Free Trial Case Manager.**

Your Case Manager will help you during the call:

- If you have not yet signed the HelloHYQVIA Form, your Case Manager will assist with that process
- Connects you with the specialty pharmacy, RxCrossroads, that will coordinate the shipment of your medication
- Connects you with a nurse from the Specialty Pharmacy Nursing Network (SPNN), who will help schedule your in-home administration training

### What you need to do:

*Keep an eye out for a call or voicemail from your Hello Case Manager. Note that this call may come from a number you do not recognize.*



## Start

A SPNN nurse will come to your home to administer your HyQvia infusions and provide initial administration training.

This will include:

- How to set up the supplies and self-infuse HyQvia
- How to optimize your HyQvia infusion rate to lessen your infusion time

### What you need to do:

*Have your Infusion Checklist (included with this brochure) ready and make sure your SPNN nurse fills in the appropriate information.*

### After the HelloHYQVIA Free Trial Program:

If you and your HCP wish for you to continue on HyQvia, you will need a prescription.

Since the insurance approval process can be lengthy, it's important that you follow up with your HCP office **prior to your last infusion** to allow enough time for this process.



### Accompanying Infusion Checklist

Use the Infusion Checklist with your SPNN nurse as you are learning how to administer HyQvia.

Keep this checklist on hand.

Should you and your HCP decide for you to continue HyQvia therapy, you will work with a different specialty pharmacy (SP) to manage your new prescription. As part of that process, your SP may provide you with a new nurse to help you formally onboard onto HyQvia as ongoing therapy.

Plan to share the Infusion Checklist with your new specialty pharmacy nurse to help with the therapy onboarding process.

## IMPORTANT SAFETY INFORMATION (continued)

### Who should not take HyQvia?

Do not take HyQvia if you:

- Are allergic to IgG, hyaluronidase, other blood products, or any ingredient in HyQvia.

### What should I avoid while taking HyQvia?

- HyQvia can make vaccines (like measles/mumps/rubella or chickenpox vaccines) not work as well for you. Before you get any vaccines, tell your HCP you take HyQvia.

### What should I tell my HCP before I start using or while using HyQvia?

Tell your HCP if you:

- Have or had any kidney, liver, or heart problems or history of blood clots because HyQvia can make these problems worse.
- Have IgA deficiency or a history of severe allergic reactions to IgG or other blood products.
- Are pregnant, trying to become pregnant or are breast feeding. It is not known whether HyQvia can harm the unborn baby or breastfed infant.

## IMPORTANT SAFETY INFORMATION (continued)

### What are the possible or reasonably likely side effects of HyQvia?

**HyQvia can cause serious side effects. If any of the following problems occur after starting HyQvia, stop the infusion immediately and contact your HCP or call emergency services:**

- Hives, swelling in the mouth or throat, itching, trouble breathing, wheezing, fainting or dizziness. These could be signs of a serious allergic reaction.
- Bad headache with nausea, vomiting, stiff neck, fever, and sensitivity to light. These could be signs of irritation and swelling of the lining around your brain.
- Reduced urination, sudden weight gain, or swelling in your legs. These could be signs of a kidney problem.
- Pain, swelling, warmth, redness, or a lump in your legs or arms, other than at the infusion site(s). These could be signs of a blood clot.
- Brown or red urine, fast heart rate, yellow skin or eyes. These could be signs of a liver or blood problem.
- Chest pain or trouble breathing, blue lips or extremities. These could be signs of a serious heart or lung problem.
- Fever over 100°F. This could be a sign of an infection.

**Please see additional safety information throughout, [click here](#) for Information for Patients, including Warning about Blood Clots, and discuss with your HCP.**

## Benefits of the HelloHYQVIA Free Trial Program

- ✓ Allows you and your physician to determine if HyQvia is right for you.
- ✓ The entire cost of your first three (3) HyQvia infusions are free, including the pump, supplies, and initial administration training. Everything will be shipped directly to your home.
- ✓ Your SPNN nurse will not only administer your HyQvia therapy, but will also work with you to optimize your therapy administration parameters.


## HyQvia Allows for Flexible Administration

- ✓ HyQvia can be infused every 3 or 4 weeks in either the stomach or thighs.
- ✓ HyQvia can be self-administered or nurse-assisted at home.
- ✓ HyQvia can be administered in your physician's office or infusion center.

## Gradual Infusion Ramp-Up

After your free trial, if you and your HCP choose for you to continue on HyQvia therapy, you will begin with your full HyQvia dose (every 3 or 4 weeks) and will not be required to do the HyQvia ramp up again.

To learn more about real patient experiences with HyQvia infusion, visit [HyQvia.com/infusing-HyQvia/what-to-expect](https://www.takeda.com/infusing-HyQvia/what-to-expect).



## IMPORTANT SAFETY INFORMATION (continued)

### What are the possible or reasonably likely side effects of HyQvia? (continued)

After HyQvia infusion a temporary, soft swelling may occur around the infusion site, which may last 1 to 3 days, due to the volume of fluid infused. The following possible side effects may occur at the site of infusion and generally go away within a few hours, and are less likely after the first few infusions.

- Mild or moderate pain
- Redness
- Swelling
- Itching

The most common side effects of HyQvia are:

- Headache
- Fatigue
- Nausea
- Fever
- Vomiting

Antibodies to the hyaluronidase component of HyQvia were formed in some patients taking HyQvia. It is not known if there is any long-term effect. In theory, these antibodies could react with your body's own hyaluronidase (PH20). PH20 is present in the male reproductive tract. So far, these antibodies have not been associated with increased or new side-effects.

**These are not all the possible side effects. Talk to your HCP about any side effect that bothers you or that does not go away.**

**You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](https://www.fda.gov/medwatch), or call 1-800-FDA-1088.**

**Please see additional safety information throughout, [click here](#) for Information for Patients, including Warning about Blood Clots, and discuss with your HCP.**

For additional information, contact HelloHYQVIA at 1.866.888.0660.

