

HyQvia

[Immune Globulin Infusion 10% (Human)
with Recombinant Human Hyaluronidase]

LEARN WHAT IT CAN MEAN

FOR YOU

HelloHYQVIA
Free Trial Program

The HelloHYQVIA Free Trial Program was designed with you in mind

With the HelloHYQVIA Free Trial Program, you can try HYQVIA for free to determine, along with your doctor, if it's right for you.*

*To be eligible, you must be a patient with an ICD-10-verified diagnosis, be a new patient not currently using HYQVIA, and have not been previously enrolled in the HelloHYQVIA program.

ICD-10=International Statistical Classification of Diseases, Tenth Revision.



Highlights of the HelloHYQVIA Free Trial Program

- ✓ **HYQVIA can be administered by a nurse at home, your physician's office, or your infusion center. Supplies will be shipped directly to your home, if applicable.**
- ✓ **Your nurse will guide you through the infusion process, either in person or virtually, to help optimize your HYQVIA infusion parameters.**
- ✓ **The entire cost of your first 3 HYQVIA infusions for PI and first 4 HYQVIA infusions for CIDP during your ramp-up period is free, including the pump, supplies, and initial infusion training.**

CIDP=chronic inflammatory demyelinating polyneuropathy; PI=primary immunodeficiency.

What is HYQVIA?

HYQVIA is a liquid medicine that is given under the skin (subcutaneously) to treat primary immunodeficiency (PI) in people 2 years and older and chronic inflammatory demyelinating polyneuropathy (CIDP) in adults.

IMPORTANT SAFETY INFORMATION

What is the most important information that I should know about HYQVIA?

- HYQVIA can cause blood clots.
- Call your healthcare professional (HCP) if you have pain, swelling, warmth, redness, or a lump in your legs or arms, other than at the infusion site(s), unexplained shortness of breath, chest pain or discomfort that worsens on deep breathing, unexplained rapid pulse, numbness or weakness on one side of the body.
- Your HCP may perform blood tests regularly to check your IgG level.
- Do not infuse HYQVIA into or around an infected or red swollen area because it can cause infection to spread.

Please see additional safety information throughout and click for [Information For Patients](#) and discuss with your HCP.

HERE'S THE PROCESS

01

Take the first step

Once you and your HCP decide to enroll in the HelloHYQVIA Free Trial Program, your HCP will verify your eligibility* and start the enrollment process.

What you need to do

If you do not have epinephrine at home already, you will need to get a prescription from your HCP. As a safety measure, epinephrine needs to be on hand for your first free trial HYQVIA infusion.

02

Meet your Case Manager

Before your first infusion, you'll receive a welcome call from a HelloHYQVIA Case Manager who will help you with the following:

- Completing your HelloHYQVIA Free Trial Form, if needed.
- Connecting with the SP, RxCrossroads, that will coordinate the shipment of your medication.
- Connecting with a nurse who will help schedule your in-home administration training.

What you need to do

Keep an eye out for a call or voicemail from your Case Manager (this call may come from a number you do not recognize).

03

Administer at home with support

A nurse will come to your home to administer your HYQVIA infusions and provide initial administration training. This will include:

- How to set up the supplies for infusion of HYQVIA.
- Detailed instructions for administration.

What you need to do

Have your HelloHYQVIA Checklist ready and make sure your nurse fills in the appropriate information.

*To be eligible you must be a patient with an ICD-10-verified diagnosis, be a new patient not currently using HYQVIA, and have not been previously enrolled in the HelloHYQVIA program.
HCP=healthcare professional; SP=specialty pharmacy.

HYQVIA is the only immune globulin subcutaneous treatment that can be:



Infused up to once a month[†]
(up to 28 days between infusions) in either the stomach or thighs.




Self-administered[‡] or nurse-assisted at home.



Administered at home[‡] or in your physician's office or infusion center.

[†]After ramp-up, every 3 or 4 weeks for PI or every 2, 3, or 4 weeks for CIDP.
[‡]After appropriate training.



HelloHYQVIA Checklist

This checklist can be used by you and your nurse to help monitor your infusions and any reactions during the free trial program.

IMPORTANT SAFETY INFORMATION (Continued)

Who should not take HYQVIA?

Do not take HYQVIA if you:

- Are allergic to IgG, hyaluronidase, other blood products, or any ingredient in HYQVIA.

What should I avoid while taking HYQVIA?

- HYQVIA can make vaccines (like measles/mumps/rubella or chickenpox vaccines) not work as well for you. Before you get any vaccines, tell your HCP that you take HYQVIA.

What should I tell my HCP before I start using or while using HYQVIA?

Tell your HCP if you:

- Have or had any kidney, liver, or heart problems or history of blood clots because HYQVIA can make these problems worse.
- Have IgA deficiency or a history of severe allergic reactions to IgG or other blood products.
- Are pregnant, trying to become pregnant or are breast feeding. It is not known whether HYQVIA can harm the unborn baby or breastfed infant.

Please see additional safety information, including Warning about Blood Clots, throughout and click for [Information For Patients and discuss with your HCP.](#)

HelloHYQVIA Free Trial Program

HyQvia
[Immune Globulin Infusion 10% (Human)
with Recombinant Human Hyaluronidase]

Infuse up to monthly*

After your free trial, if you and your HCP decide HYQVIA is right for you, you will need a prescription. The insurance approval process can be lengthy. **Reach out to your HCP's office prior to your last infusion provided by the free trial program to allow enough time for this process.** Thereafter, you will continue with HYQVIA based on your doctor's dosing schedule (every 2, 3, or 4 weeks).* You will also work with a different SP to manage your new prescription. As part of that process, your SP may provide you with a new nurse to help you formally onboard onto HYQVIA as ongoing therapy.

Click here to learn more about patient experiences with HYQVIA.

*After ramp-up, every 3 or 4 weeks for PI or every 2, 3, or 4 weeks for CIDP.

Start the conversation about the HelloHYQVIA program

If you're eligible, the HelloHYQVIA program lets you try HYQVIA for free to see, along with your doctor, if it's right for you

For additional information or to learn how to enroll in the HelloHYQVIA Free Trial Program, contact your HCP, visit HYQVIA.com, or call **1-866-861-1750**.

IMPORTANT SAFETY INFORMATION (Continued)

What are the possible or reasonably likely side effects of HYQVIA?

HYQVIA can cause serious side effects. If any of the following problems occur after starting HYQVIA, stop the infusion immediately and contact your HCP or call emergency services:

- Hives, swelling in the mouth or throat, itching, trouble breathing, wheezing, fainting or dizziness. These could be signs of a serious allergic reaction.
- Bad headache with nausea, vomiting, stiff neck, fever, and sensitivity to light. These could be signs of irritation and swelling of the lining around your brain.
- Reduced urination, sudden weight gain, or swelling in your legs. These could be signs of a kidney problem.
- Pain, swelling, warmth, redness, or a lump in your legs or arms, other than at the infusion site(s). These could be signs of a blood clot.
- Brown or red urine, fast heart rate, yellow skin or eyes. These could be signs of a liver or blood problem.
- Chest pain or trouble breathing, blue lips or extremities. These could be signs of a serious heart or lung problem.
- Fever over 100°F. This could be a sign of an infection.

After HYQVIA infusion a temporary, soft swelling may occur around the infusion site, which may last 1 to 3 days, due to the volume of fluid infused. The following possible side effects may occur at the site of infusion and generally go away within a few hours, and are less likely after the first few infusions.

- Mild or moderate pain
- Redness
- Swelling
- Itching

The most common side effects of HYQVIA in PI are:

- Headache
- Fatigue
- Nausea
- Fever
- Vomiting

The most common side effects of HYQVIA in CIDP are:

- Headache
- Fatigue
- Nausea
- Fever
- Itching
- Redness
- Abdominal pain
- Back pain
- Pain in extremity

Antibodies to the hyaluronidase component of HYQVIA were formed in some patients taking HYQVIA. It is not known if there is any long-term effect. In theory, these antibodies could react with your body's own hyaluronidase (PH20). PH20 is present in the male reproductive tract. So far, these antibodies have not been associated with increased or new side-effects.

These are not all the possible side effects. Talk to your HCP about any side effect that bothers you or that does not go away.

For additional safety information, including Warning about Blood Clots, click for [Information For Patients](#) and discuss with your HCP.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call **1-800-FDA-1088.**

